

Long-range order and spin liquid states of polycrystalline $\text{Tb}_{2+x}\text{Ti}_{2-x}\text{O}_{7+y}$

T. Taniguchi,¹ H. Kadowaki,¹ H. Takatsu,¹ B. Fåk,² J. Ollivier,³ T. Yamazaki,⁴ T. J. Sato,⁵ H. Yoshizawa,⁵ Y. Shimura,⁴ T. Sakakibara,⁴ T. Hong,⁶ K. Goto,¹ L. R. Yaraskavitch,⁷ and J. B. Kycia⁷

¹*Department of Physics, Tokyo Metropolitan University, Hachioji-shi, Tokyo 192-0397, Japan*

²*SPSMS, UMR-E CEA/UJF-Grenoble-1, INAC, F-38054 Grenoble Cedex 9, France*

³*Institute Laue Langevin, BP156, F-38042 Grenoble, France*

⁴*Institute for Solid State Physics, University of Tokyo, Kashiwa 277-8581, Japan*

⁵*NSL, Institute for Solid State Physics, University of Tokyo, Tokai, Ibaraki 319-1106*

⁶*Quantum Condensed Matter Division, Oak Ridge National Laboratory, Oak Ridge, TN 37831-6393, USA*

⁷*Department of Physics and Astronomy and Guelph-Waterloo Physics Institute, University of Waterloo, Waterloo, ON N2L 3G1, Canada*

(Dated: November 16, 2012)

Low-temperature states of polycrystalline samples of a frustrated pyrochlore oxide $\text{Tb}_{2+x}\text{Ti}_{2-x}\text{O}_{7+y}$ have been investigated by specific heat, magnetic susceptibility, and neutron scattering experiments. We have found that this system can be tuned from a long-range ordered state ($x > x_c$) to a spin-liquid state by a minute change of x . Specific heat shows a sharp peak at a phase transition at $T_c = 0.5$ K for $x = 0.005$. Inelastic neutron scattering shows that the crystal field ground state doublet of Tb^{3+} splits into two singlets below T_c , suggesting a cooperative Jahn-Teller transition due to a magneto-elastic coupling, accompanied by a small antiferromagnetic ordering.

PACS numbers: 75.10.Kt, 75.40.Cx, 75.70.Tj, 78.70.Nx

Magnetic systems with geometric frustration, a prototype of which is antiferromagnetically coupled Ising spins on a triangle, have been intensively studied experimentally and theoretically for decades¹. Spin systems on networks of triangles or tetrahedra, such as triangular², kagomé³, and pyrochlore⁴ lattices, play major roles in these studies. Subjects that have fascinated many investigators in recent years are classical and quantum spin-liquid states^{5–8}, where conventional long-range order (LRO) is suppressed to very low temperatures. Quantum spin-liquids^{6,7} in particular have been challenging both theoretically and experimentally since the proposal of the resonating valence-bond state⁹. The spin ice materials $\text{R}_2\text{Ti}_2\text{O}_7$ ($\text{R} = \text{Dy}, \text{Ho}$) are the well-known classical examples⁵, while other experimental candidates found recently have been studied^{10–12}.

Among frustrated pyrochlore oxides⁴, $\text{Tb}_2\text{Ti}_2\text{O}_7$ has attracted much attention because it does not show any conventional LRO down to 50 mK and remains in a dynamic spin-liquid state^{13,14}. Theoretical considerations of the crystal-field (CF) states of Tb^{3+} and exchange and dipolar interactions of the system^{15–17} showed that it should undergo a transition into a magnetic LRO state at about 1.8 K within a random phase approximation¹⁷. The puzzling origin of the spin-liquid state of $\text{Tb}_2\text{Ti}_2\text{O}_7$ is in hot debate^{4,18–25}. An interesting scenario to explain the spin-liquid state is the theoretical proposal of a quantum spin-ice state¹⁹. More recently, another scenario of a two-singlet spin-liquid state was proposed to explain why inelastic neutron spectra in a low energy range are observed despite the fact Tb^{3+} is a non-Kramers ion^{20,21}.

Several experimental puzzles of $\text{Tb}_2\text{Ti}_2\text{O}_7$ originate from the difficulty of controlling the quality of single crystalline samples, resulting in strongly sample-dependent specific-heat anomalies at temperatures below

2 K^{15,23,26–29}. In contrast, experimental results on polycrystalline samples are more consistent^{13,14,23}. Among experimental results reported to date, an important clue to solve the puzzles seems to be a change of state at about 0.4 K suggested by specific heat²³, inelastic neutron scattering²³, and neutron spin echo¹⁴ on polycrystalline samples. At this temperature, a few single-crystalline samples show a peak in the specific heat suggesting a phase transition^{26,27}, an issue that has not been pursued seriously. A possibility of a cooperative Jahn-Teller phase-transition well below 1 K was inferred many years ago from the observation of an anomalous temperature dependence of the elastic constants above 1 K³⁰. The two-singlet spin-liquid scenario of Refs. 20, 21, and 31 is based on the assumption of a tetragonal lattice distortion in $\text{Tb}_2\text{Ti}_2\text{O}_7$ and the closely related ordered spin-ice compound $\text{Tb}_2\text{Sn}_2\text{O}_7$ ³², but the accompanying lattice distortion might be too difficult to observe directly^{22,33–36}.

In the present work, we investigate the hypothesis that the non-stoichiometry x of $\text{Tb}_{2+x}\text{Ti}_{2-x}\text{O}_{7+y}$ is a tuning parameter for a quantum critical point separating a LRO state from a spin liquid state. We have therefore performed specific heat, magnetization, and neutron scattering experiments on polycrystalline samples of $\text{Tb}_{2+x}\text{Ti}_{2-x}\text{O}_{7+y}$ with different values of x . We find that a minute change of x brings about a systematic change of the specific heat. The ground state goes from LRO for $x > x_c$ to a spin liquid for $x < x_c$. Inelastic neutron scattering strongly suggests that this LRO is a cooperative Jahn-Teller lattice distortion accompanied by a small antiferromagnetic ordering. If this interpretation is correct, we may make a conjecture that the ground state for $x < x_c$ is a spin liquid state in which spin and lattice degrees of freedom are governed by quantum fluctuations.

Polycrystalline samples of $\text{Tb}_{2+x}\text{Ti}_{2-x}\text{O}_{7+y}$ with




FIG. 1. Lattice constants of polycrystalline $\text{Tb}_{2+x}\text{Ti}_{2-x}\text{O}_{7+y}$ at 25 °C. The dashed line is a guide to the eye.

$-0.015 < x < 0.01$ were prepared by standard solid-state reaction¹³. The value of x was adjusted by changing the mass ratio of the two starting materials, Tb_4O_7 and TiO_2 , which were heated in air at 1350 °C for several days with periodic grindings to ensure a complete reaction. It was ground into powder and annealed in air at 800°C for one day. The values of x used in this report are nominal, and have an offset about ± 0.002 . The value of y is determined by the oxidizing conditions. X-ray powder-diffraction experiments were carried out using a RIGAKU-SmartLab powder diffractometer equipped with a Cu $K_{\alpha 1}$ monochromator. The absence of impurity peaks in the powder diffraction patterns shows that the samples are single phase with pyrochlore structure³⁷. To measure the x dependence of the lattice constant a at 25 °C, we performed θ -2 θ scans on powder mixtures of $\text{Tb}_{2+x}\text{Ti}_{2-x}\text{O}_{7+y}$ and Si. Figure 1 shows that the lattice constant a has a smooth variation with x , which ensures a continuous change of the stoichiometry of $\text{Tb}_{2+x}\text{Ti}_{2-x}\text{O}_{7+y}$ for small x .

Specific heat above 0.4 K was measured on a physical-property measurement-system. Measurements below 0.4 K were carried out using the quasi-adiabatic relaxation method on a dilution refrigerator³⁸. DC magnetization measurements were carried out by a capacitive Faraday magnetometer in a ^3He refrigerator. Neutron powder diffraction measurements were performed on the triple-axis spectrometer CTAX at ORNL. Inelastic neutron scattering measurements were carried out on the time-of-flight spectrometer IN5 operated with $\lambda = 5$ and 10 Å at ILL. For these neutron scattering experiments, samples of $x = 0.005$ and -0.005 with weights of 5 and 9 g were mounted in a ^3He (CTAX) and a dilution refrigerator (IN5), respectively.

In Fig. 2 we show the specific heat C_P of the polycrystalline samples as a function of temperature together with a few previous measurements^{23,26,39}. Earlier work have shown qualitatively similar results^{40,41}. The $C_P(T)$




FIG. 2. (Color online) Temperature dependence of the specific heat of polycrystalline $\text{Tb}_{2+x}\text{Ti}_{2-x}\text{O}_{7+y}$. Previous measurements of poly- and single-crystalline samples^{23,26,39}, as well as the present measurements below 0.6 K of a sample prepared in the same manner as in Ref. 23, are plotted for comparison. The inset shows a phase diagram expected from the specific heat, susceptibility, and neutron scattering.

data show a systematic change by varying x . A sample with $x = 0.005$ shows a clear peak indicating a second-order phase transition at $T_c = 0.5$ K. Samples with $x = 0.0025$ and 0.000 show smaller peaks at 0.43 and 0.4 K, respectively. We note that C_P of the present sample with $x = 0.000$ agrees approximately with our previous measurements²³, the temperature range of which was extended down to 0.2 K in the present work on a sample (nominal $x' = 0$) prepared from a different commercial source of Tb_4O_7 . Our previous interpretation²³ of the upturn below 0.5 K as a crossover behavior is incorrect owing to the insufficient temperature range. The previous C_P data³⁹ (Fig. 2) on a polycrystalline sample with their nominal $x'' = 0$ corresponds to our $x = -0.0125$, implying that fine tuning of x requires careful sample preparation. In the inset of Fig. 2, we show a cumulative phase diagram constructed from $C_P(T, x)$ in conjunction with the susceptibility and neutron scattering experiments discussed below.

A peak of $C_P(T)$ in $\text{Tb}_2\text{Ti}_2\text{O}_7$ was first reported for a single-crystalline sample at 0.37 K²⁶. These $C_P(T)$ data²⁶, reproduced in Fig. 2, show significantly different T dependence from any of the polycrystalline samples. The sharp peak at 0.37 K may result from a portion of




FIG. 3. (Color online) Temperature dependence of the magnetic susceptibility of polycrystalline $\text{Tb}_{2+x}\text{Ti}_{2-x}\text{O}_{7+y}$ with $x = -0.005, 0.000$, and 0.005 .

the sample having a non-stoichiometry parameter around $x = -0.001$, corresponding to a peak slightly lower in temperature than our $x = 0.000$. However, a hump in $C_P(T)$ around 0.75 K for the single crystal does not appear for the polycrystalline samples. We believe that these single- and poly-crystalline samples have significant, but presently not well understood, differences in quality.

In order to check whether T_c is an antiferromagnetic transition, as suggested in Ref. 26, we performed magnetization and neutron powder-diffraction experiments. In Fig. 3 we show the magnetic susceptibility as a function of temperature for three polycrystalline samples with $x = \pm 0.005$ and 0.000 . The susceptibilities for $x = 0.005$ and 0.000 show only slight anomalies around the clear peaks of $C_P(T)$ at $T_c = 0.5$ and 0.4 K, respectively. These results are very different from typical behavior expected of antiferromagnetic phase transitions.

In Fig. 4 we show neutron powder-diffraction patterns for the $x = 0.005$ sample below and above T_c . The pattern below T_c shows neither any clear antiferromagnetic reflections nor any clear changes due to a structural transition. A rough estimate of the upper limit of the antiferromagnetic ordered moment is about $0.1 \mu_B$. The intensity of the sloping paramagnetic scattering, a background for Bragg peaks, decreases slightly as temperature is lowered from 1.2 to 0.28 K. This is brought about by a change in the magnetic excitations. The lack of obvious antiferromagnetism distinctly separates $\text{Tb}_2\text{Ti}_2\text{O}_7$ from the ordered spin-ice compound $\text{Tb}_2\text{Sn}_2\text{O}_7$ ^{32,42}, in which antiferromagnetic ordering with a moment of $5.9 \mu_B$ was observed well below $T_c = 0.87$ K.

To study the spectral change of the magnetic excitations through T_c , we performed inelastic neutron scattering measurements using the spectrometer IN5⁴³ with an energy resolution of $\Delta E = 0.01$ meV (FWHM), which is 6 times better than in a previous study²³. Figure 5 shows




FIG. 4. (Color online) Neutron powder diffraction pattern of polycrystalline $\text{Tb}_{2+x}\text{Ti}_{2-x}\text{O}_{7+y}$ with $x = 0.005$ taken above and below $T_c = 0.5$ K.

the temperature dependence of an energy spectrum for the $x = 0.005$ sample at $Q = 0.6 \text{ \AA}^{-1}$. It is evident that the spectrum changes from a continuum ($T > T_c$) to a peaked structure at 0.1 meV ($T < T_c$). Since the peak at 0.1 meV is essentially dispersionless and Q independent, the excitation is probably due to a CF splitting of the ground state doublet by a lowering of the local trigonal symmetry. As pointed out in Refs. 20, 21, and 31, a splitting of the CF ground-state doublet into two singlets is the simplest evidence of a Jahn-Teller distortion due to a magneto-elastic coupling. Therefore, the evolution of the 0.1 meV excitation strongly suggests that T_c is a Jahn-Teller structural phase transition inferred in Ref. 30. An energy spectrum of the $x = -0.005$ sample is also shown in Fig. 5 for comparison, revealing quantum fluctuations with an energy scale of 0.1 meV.

The high sensitivity of IN5 enabled us to observe a small Bragg peak, being undetectable in the CTAX data (Fig. 4). In the inset of Fig. 5, the intensity of the elastic scattering for $|E| < 0.005$ meV is plotted as a function of Q . Below T_c , a clear Bragg peak at $Q = 0.54 \text{ \AA}^{-1}$ is observed, which can be indexed as $(\frac{1}{2} \frac{1}{2} \frac{1}{2})$. Although this peak could be of a nuclear (structural) origin, it is more likely an antiferromagnetic reflection. In fact, two recent neutron scattering experiments carried out on single-crystalline samples of $\text{Tb}_2\text{Ti}_2\text{O}_7$ show magnetic short-range-order around the same $Q = (\frac{1}{2} \frac{1}{2} \frac{1}{2})$ ^{21,25}. A rough estimate of the antiferromagnetic ordered moment is $0.08 \mu_B$. We note that the Q -width of the $(\frac{1}{2} \frac{1}{2} \frac{1}{2})$ peak is somewhat larger than the coarse Q resolution on IN5. Whether this peak is truly long-range order would have to await high-resolution neutron diffraction measurements, which are difficult in view of the small moment.

The previously reported transition or crossover at about 0.4 K for the poly- and single-crystalline




FIG. 5. (Color online) Energy spectra of inelastic neutron scattering for polycrystalline $\text{Tb}_{2+x}\text{Ti}_{2-x}\text{O}_{7+y}$ with $x = 0.005$ and -0.005 . The inset shows the Q dependence of the elastic scattering for the $x = 0.005$ sample around $Q = |\frac{1}{2}\frac{1}{2}\frac{1}{2}|$ above and below T_c .

$\text{Tb}_2\text{Ti}_2\text{O}_7$ ^{14,23,26} is presumably attributable to the same origin as that of the present $x = 0.005$ sample. The clear 0.1 meV excitation peak for this sample is most simply accounted for by a Jahn-Teller distortion and resultant CF splitting. Since this is an indirect evidence for the structural transition, more direct observation, by X-ray diffraction e.g., remains to be performed, but such measurements at low temperatures are exceedingly difficult. By assuming a Jahn-Teller distortion ($T \ll T_c$), an expansion of the theoretical framework of Refs. 20, 21, and

31 may be a promising direction to explain the ground state of the polycrystalline $\text{Tb}_{2+x}\text{Ti}_{2-x}\text{O}_{7+y}$ with $x > x_c$, especially for the analysis of inelastic neutron scattering data. In order to reproduce the phase transition at $T_c > 0$, the theory^{20,21,31} will have to be modified to include a soft phonon mode and a spin-lattice coupling. Along this line, the long-standing puzzle of the spin liquid state of $\text{Tb}_2\text{Ti}_2\text{O}_7$ may be reformulated to a novel problem of frustration having both spin and lattice degrees of freedom; Why and how do the spins and the soft phonon modes fluctuate quantum mechanically down to $T = 0$ for $x < x_c$?

In summary, we have investigated the low-temperature states of polycrystalline $\text{Tb}_{2+x}\text{Ti}_{2-x}\text{O}_{7+y}$ samples by specific heat, magnetic susceptibility, and neutron scattering experiments. We have found that this system can be tuned by a minute change of the parameter x from a LRO ground state for $x > x_c$ to a liquid-type ground state with spin- and possibly lattice-fluctuations for $x < x_c$. Specific heat shows a sharp peak at a second-order phase-transition T_c for $x > x_c$. Inelastic neutron scattering shows that the CF ground doublet splits into two singlets below T_c , suggesting strongly that T_c is a co-operative Jahn-Teller structural transition accompanied by a small antiferromagnetic ordering with a wave-vector $(\frac{1}{2}\frac{1}{2}\frac{1}{2})$.

We thank M.J.P. Gingras, R. Higashinaka, J.W. Lynn, and K. Matsuhira for useful discussions. This work was supported by KAKENHI NSMIF. The specific heat to 0.4 K and magnetization measurements were performed using facilities of ISSP, Univ. of Tokyo. Work on CTAX was supported by the US-Japan Cooperative Program on Neutron Scattering. HFIR was partially supported by the US DOE, Office of BES, Division of Scientific User Facilities. The neutron scattering performed using IN5 (France) was transferred from JRR3-HER (proposal 11567) with the approval of ISSP, Univ. of Tokyo, and JAEA, Tokai, Japan.

- ¹ C. Lacroix, P. Mendels, and F. Mila, eds., *Introduction to Frustrated Magnetism* (Springer, 2011).
- ² G. H. Wannier, Phys. Rev. **79**, 357 (1950).
- ³ I. Syôzi: Prog. Theor. Phys. **6** (1951) 306.
- ⁴ J. S. Gardner, M. J. P. Gingras, and J. E. Greedan, Rev. Mod. Phys. **82**, 53 (2010).
- ⁵ S. T. Bramwell and M. J. P. Gingras, Science **294**, 1495 (2001).
- ⁶ P. A. Lee, Science **321**, 1306 (2008).
- ⁷ L. Balents, Nature **464**, 199 (2010).
- ⁸ S. Yan, D. A. Huse, and S. R. White, Science **332**, 1173 (2011).
- ⁹ P. W. Anderson, Mater. Res. Bull. **8**, 153 (1973).
- ¹⁰ J. S. Helton, K. Matan, M. P. Shores, E. A. Nytko, B. M. Bartlett, Y. Yoshida, Y. Takano, A. Suslov, Y. Qiu, J.-H. Chung, D. G. Nocera, and Y. S. Lee, Phys. Rev. Lett. **98**, 107204 (2007).
- ¹¹ T. Itou, A. Oyamada, S. Maegawa, M. Tamura, and

- R. Kato, Phys. Rev. B **77**, 104413 (2008).
- ¹² B. Fåk, E. Kermarrec, L. Messio, B. Bernu, C. Lhuillier, F. Bert, P. Mendels, B. Koteswararao, F. Bouquet, J. Olivier, A. D. Hillier, A. Amato, R. H. Colman, and A. S. Wills, Phys. Rev. Lett. **109**, 037208 (2012).
- ¹³ J. S. Gardner, S. R. Dunsiger, B. D. Gaulin, M. J. P. Gingras, J. E. Greedan, R. F. Kiefl, M. D. Lumsden, W. A. MacFarlane, N. P. Raju, J. E. Sonier, I. Swainson, and Z. Tun, Phys. Rev. Lett. **82**, 1012 (1999).
- ¹⁴ J. S. Gardner, A. Keren, G. Ehlers, C. Stock, E. Segal, J. M. Roper, B. Fåk, M. B. Stone, P. R. Hammar, D. H. Reich, and B. D. Gaulin, Phys. Rev. B **68**, 180401 (2003).
- ¹⁵ M. J. P. Gingras, B. C. den Hertog, M. Faucher, J. S. Gardner, S. R. Dunsiger, L. J. Chang, B. D. Gaulin, N. P. Raju, and J. E. Greedan, Phys. Rev. B **62**, 6496 (2000).
- ¹⁶ M. Enjalran and M. J. P. Gingras, Phys. Rev. B **70**, 174426 (2004).
- ¹⁷ Y.-J. Kao, M. Enjalran, A. Del Maestro, H. R. Molavian,

- and M. J. P. Gingras, Phys. Rev. B **68**, 172407 (2003).
- ¹⁸ Y. Yasui, M. Kanada, M. Ito, H. Harashina, M. Sato, H. Okumura, K. Kakurai, and H. Kadowaki, J. Phys. Soc. Jpn. **71**, 599 (2002).
 - ¹⁹ H. R. Molavian, M. J. P. Gingras, and B. Canals, Phys. Rev. Lett. **98**, 157204 (2007).
 - ²⁰ P. Bonville, I. Mirebeau, A. Gukasov, S. Petit, and J. Robert, Phys. Rev. B **84**, 184409 (2011).
 - ²¹ S. Petit, P. Bonville, J. Robert, C. Decorse, and I. Mirebeau, Phys. Rev. B **86**, 174403 (2012).
 - ²² B. D. Gaulin, J. S. Gardner, P. A. McClarty, and M. J. P. Gingras, Phys. Rev. B **84**, 140402 (2011).
 - ²³ H. Takatsu, H. Kadowaki, T. J. Sato, J. W. Lynn, Y. Tabata, T. Yamazaki, and K. Matsuhira, J. Phys.: Condens. Matter **24**, 052201 (2012).
 - ²⁴ T. Fennell, M. Kenzelmann, B. Roessli, M. K. Haas, and R. J. Cava, Phys. Rev. Lett. **109**, 017201 (2012).
 - ²⁵ K. Fritsch, K. A. Ross, Y. Qiu, J. R. D. Copley, T. Guidi, R. I. Bewley, H. A. Dabkowska, and B. D. Gaulin, arXiv:1210.1242.
 - ²⁶ N. Hamaguchi, T. Matsushita, N. Wada, Y. Yasui, and M. Sato, Phys. Rev. B **69**, 132413 (2004).
 - ²⁷ Y. Chapuis, Ph.D. thesis, Université Joseph Fourier (2009), <http://tel.archives-ouvertes.fr/tel-00463643/en/>.
 - ²⁸ Y. Chapuis, A. Yaouanc, P. Dalmas de Réotier, C. Marin, S. Vanishri, S. H. Curnoe, C. Vâju, and A. Forget, Phys. Rev. B **82**, 100402 (2010).
 - ²⁹ A. Yaouanc, P. Dalmas de Réotier, Y. Chapuis, C. Marin, S. Vanishri, D. Aoki, B. Fåk, L.-P. Regnault, C. Buisson, A. Amato, C. Baines, and A. D. Hillier, Phys. Rev. B **84**, 184403 (2011).
 - ³⁰ L. G. Mamsurova, K. S. Pigal'skiĭ, and K. K. Pukhov, JETP Lett. **43**, 755 (1986).
 - ³¹ S. Petit, P. Bonville, I. Mirebeau, H. Mutka, and J. Robert, Phys. Rev. B **85**, 054428 (2012).
 - ³² I. Mirebeau, A. Apetrei, J. Rodríguez-Carvajal, P. Bonville, A. Forget, D. Colson, V. Glazkov, J. P. Sanchez, O. Isnard, and E. Suard, Phys. Rev. Lett. **94**, 246402 (2005).
 - ³³ T. T. A. Lummen, I. P. Handayani, M. C. Donker, D. Fausti, G. Dhalenne, P. Berthet, A. Revcolevschi, and P. H. M. van Loosdrecht, Phys. Rev. B **77**, 214310 (2008).
 - ³⁴ J. P. C. Ruff, B. D. Gaulin, J. P. Castellan, K. C. Rule, J. P. Clancy, J. Rodriguez, and H. A. Dabkowska, Phys. Rev. Lett. **99**, 237202 (2007).
 - ³⁵ Y. Nakanishi, T. Kumagai, M. Yoshizawa, K. Matsuhira, S. Takagi, and Z. Hiroi, Phys. Rev. B **83**, 184434 (2011).
 - ³⁶ K. Goto, H. Takatsu, T. Taniguchi, and H. Kadowaki, J. Phys. Soc. Jpn. **81**, 015001 (2012).
 - ³⁷ S.-W. Han, J. S. Gardner, and C. H. Booth, Phys. Rev. B **69**, 024416 (2004).
 - ³⁸ J. A. Quilliam, C. G. A. Mugford, A. Gomez, S. W. Kycia, and J. B. Kycia, Phys. Rev. Lett. **98**, 037203 (2007).
 - ³⁹ R. Siddharthan, B. S. Shastry, A. P. Ramirez, A. Hayashi, R. J. Cava, and S. Rosenkranz, Phys. Rev. Lett. **83**, 1854 (1999).
 - ⁴⁰ A. Cornelius, B. Light, R. S. Kumar, M. Eichenfield, T. Dutton, R. Pepin, and J. Gardner, Physica B: Condensed Matter **359-361**, 1243 (2005).
 - ⁴¹ X. Ke, D. V. West, R. J. Cava, and P. Schiffer, Phys. Rev. B **80**, 144426 (2009).
 - ⁴² I. Mirebeau, P. Bonville, and M. Hennion, Phys. Rev. B **76**, 184436 (2007).
 - ⁴³ J. Ollivier and H. Mutka, J. Phys. Soc. Jpn. **80**, SB003 (2011).